

### Catrgories and functors

**Notations.** We write  $Set, Top, Ab, Grp, Cmr, Mod_K, Vec_{\mathbb{k}} = Mod_{\mathbb{k}}, Ass_{\mathbb{k}}, A-Mod, Mod-A$  for the categories of sets, topological spaces, abelian groups, all groups, commutative rings<sup>1</sup>, modules over commutative ring  $K$ , vector spaces and associative algebras over a field  $\mathbb{k}$ , left and right modules over algebra  $A$  respectively. Categories of functors  $\mathcal{C} \rightarrow \mathcal{D}$  and presheaves<sup>2</sup>  $\mathcal{C}^{opp} \rightarrow \mathcal{D}$  are denoted by  $Fun(\mathcal{C}, \mathcal{D})$  and  $pSh(\mathcal{C}, \mathcal{D})$ .

**SHA1◊1.** Let  $\Delta_{big}$  be the category of all finite ordered sets with order preserving maps as the morphisms and  $\Delta \subset \Delta_{big}$  be its full small subcategory formed by sets  $[n] \stackrel{def}{=} \{0, 1, \dots, n\}, n \geq 0$ , ordered usually. Show that a)  $\Delta$  and  $\Delta_{big}$  are equivalent b) algebra  $\mathbb{Z}[\Delta]$  is generated by the identity arrows  $e_n = Id_{[n]}$ , the inclusions  $\partial_n^{(i)} : [n-1] \hookrightarrow [n], 0 \leq i \leq n, i \notin \partial_n^{(i)}([n-1])$ , and surjections  $s_n^{(i)} : [n] \twoheadrightarrow [n-1], 0 \leq i \leq n-1, (i+1) \mapsto i$ . **в\*)** Find generators for the ideal of relations between these generating arrows.

**SHA1◊2.** For a given  $X \in Ob \mathcal{C}$  let a functor  $h^X : Y \mapsto Hom(X, Y)$  and a presheaf  $h_X : Y \mapsto Hom(Y, X)$  take an arrow  $\varphi : Y_1 \rightarrow Y_2$  respectively to the left and right multiplications by this arrow:

$$\varphi_* : Hom(X, Y_1) \rightarrow Hom(X, Y_2), \psi \mapsto \varphi \circ \psi \quad \text{and} \quad Hom(Y_2, X) \rightarrow Hom(Y_1, X), \psi \mapsto \psi \circ \varphi.$$

Show that prescriptions  $X \mapsto h^X$  and  $X \mapsto h_X$  define a pre-sheaf  $h^* : \mathcal{C}^{opp} \rightarrow Fun(\mathcal{C}, Set)$  and a functor  $h_* : \mathcal{C} \rightarrow pSh(\mathcal{C}, Set)$  respectively.

**SHA1◊3.** Show that functor  $h^X : Ab \rightarrow Ab$  takes an exact sequence  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  to an exact sequence  $0 \rightarrow Hom(X, A) \rightarrow Hom(X, B) \rightarrow Hom(X, C)$  whose rightmost arrow may be non-surjective. Formulate and prove dual property of functor  $h_X : Ab \rightarrow Ab$ .

**SHA1◊4.** Describe products and coproducts in a)  $Set$  б)  $Top$  в)  $Mod_K$  г)  $Grp$  д)  $Cmr$ .

**SHA1◊5.** Fix prime  $p \in \mathbb{N}$ . For each  $n \in \mathbb{N}$  let  $A_n = \mathbb{Z}/(p^n)$ . For  $m > n$  write  $\psi_{nm} : A_m \twoheadrightarrow A_n$  for the factorization mapping and  $\varphi_{mn} : A_n \hookrightarrow A_m$  for the embedding  $[1] \mapsto [p^{m-n}]$ . In category  $Ab$  describe a)  $\varprojlim A_n$  along  $\psi_{mn}$  б)  $\varinjlim A_n$  along  $\varphi_{mn}$ .

**SHA1◊6.** Let  $B_n = \mathbb{Z}/(n)$ . For  $n|m$  write  $\psi_{nm} : B_m \twoheadrightarrow B_n$  and  $\varphi_{mn} : B_n \hookrightarrow B_m$  for the factorization mapping and the embedding  $[1] \mapsto [m/n]$  respectively. In category  $Ab$  describe a)  $\varprojlim B_n$  along  $\psi_{nm}$  б)  $\varinjlim B_n$  along  $\varphi_{mn}$ .

**SHA1◊7.** Prove that a functor  $G : \mathcal{D} \rightarrow \mathcal{C}$  admits a left adjoint functor  $F$  iff for each  $X \in Ob \mathcal{C}$  a functor  $h_G^X : Y \mapsto Hom_{\mathcal{C}}(X, G(Y))$  is corepresentable, and in this case  $F(X)$  corepresents  $h_G^X$ . Formulate and prove the dual criteria for the existence of right adjoint functor  $G$  to a given functor  $F : \mathcal{C} \rightarrow \mathcal{D}$ .

**SHA1◊8.** Show that any left adjoint functor commutes with colimits and any right adjoint functor commutes with limits<sup>3</sup>.

**SHA1◊9.** For an arbitrary extension  $S \subset R$  of associative algebras with units construct left and right adjoint functors to the restriction functor  $res_S^R : R-Mod \rightarrow S-Mod$ .

**SHA1◊10.** Given a topological space  $\mathcal{X}$ , write  $S(\mathcal{X}) : \Delta^{opp} \rightarrow Set$  for the simplicial set that takes  $[n] \in Ob \Delta$  to  $S_n(\mathcal{X}) \stackrel{def}{=} Hom_{Top}(\Delta^n, \mathcal{X})$ , where  $\Delta^n \subset \mathbb{R}^{n+1}$  is the standard regular  $n$ -dimensional simplex, and takes an order preserving arrow  $\varphi : [n] \rightarrow [m]$  to the right multiplication mapping  $f \mapsto f \circ |\varphi|$ , where  $|\varphi| : \Delta^n \rightarrow \Delta^m$  stays for the affine mapping acting on the vertices as  $\varphi$ . Show that the functor  $S : Top \rightarrow pSh(\Delta)$  is right adjoint to the geometric realization functor  $pSh(\Delta) \rightarrow Top$ .

<sup>1</sup>with unity and homomorphisms sending unity to unity

<sup>2</sup>i.e. contravariant functors

<sup>3</sup>functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  commutes with (co)limits, if for each  $L \in Ob \mathcal{C}$  and any diagram  $\Phi : \mathcal{N} \rightarrow \mathcal{C}$  the condition « $L$  is the (co)limit of  $\Phi$  in  $\mathcal{C}$ » implies the condition « $F(L)$  is the (co)limit of  $F \circ \Phi$  in  $\mathcal{D}$ »

Персональный табель \_\_\_\_\_  
(напишите свои имя, отчество и фамилию)

Task № 1 (September 4, 2015)

№	дата сдачи	имя и фамилия принявшего	подпись принявшего
1а			
б			
в			
2			
3			
4а			
б			
в			
г			
д			
5а			
б			
6а			
б			
7			
8			
9			
10			